Intersatellite Calibrated HIRS Upper Tropospheric Water Vapor

Lei Shi and John J. Bates

NOAA National Climatic Data Center Asheville, North Carolina, U.S.A.





NOAA National Climatic Data Center

Outline

- Motivation
 - Time series discontinuity from satellite to satellite, particularly from TOVS to ATOVS
 - Upper tropospheric water vapor (UTWV) is an important fundamental climate data record (CDR)
 - UTWV is key to water vapor feedback
- Approach
 - New and improved intersatellite calibration based on two datasets
 - Overlaps of zonal means
 - Simultaneous nadir overpass (SNO)
- Result
 - Extended time series of the fundamental CDR to present





Motivation – Uncorrected Intersatellite Differences of UTWV (Channel 12)



- Due to the independence of individual HIRS instrument's calibration, biases exist from satellite to satellite.
- These intersatellite biases have become a common source of uncertainty faced by long-term studies.





Spectral Filter Functions



- Differences between HIRS/2 and HIRS/3 are expected due to different filter functions.
- In-orbit performance still has biases unexplained by filter functions.
 - Thus empirical approach is considered.





Approach – Two Datasets for Intersatellite Calibration to Cover Diverse Atmospheres

- Overlaps of zonal means for tropical and midlatitude atmosphere cases
 - Zonal means of channel 12 (UTWV) are computed for every 10-degree latitude belt.
 - Differences are derived from overlapping satellites.
- Simultaneous nadir overpass (SNO) observations for polar cases
 - SNOs occur when two satellites cross each other
 - Data taken at the same location at the satellite nadir within a few seconds

7-13 May 2008

- In the regions 70-80N and 70-80S





Temperature-dependent Intersatellite Differences from Zonal Mean Approach





Temperature-dependent Intersatellite Differences from SNO Approach



Biases are also temperature dependent.



Extension of Time Series

- Orbital, daily, and monthly, 1979-present
- At both pixel resolution and 2.5x2.5 lat/lon grids
- Intersatellitecalibrated



30S - 30N

* V1 is derived based on methods discussed in Bates, J. J., X. Wu, and D. L. Jackson, (1996) Interannual variability of upper-troposphere water vapor band brightness temperatures. J. Climate, **9**, 428-438.



Intersatellite Calibrated to N-12 (showing 30S - 30N)

- **Biases** minimized.
- Temperature dependent biases accounted for.
- Similar overall variances between HIRS/2 and HIRS/3.
- Time series can be extended as variance preserved.





Climatologic Mean for Jan, Apr, Jul, and Oct









Seasonal Variation

- Within tropics, drier atmosphere produces a higher value of UTWV (channel 12 "sees" lower atmosphere).
- Minima in the tropics indicate moist upper troposphere caused by convection.
- Maxima in northern tropics in January move to southern tropics in July, indicating the move of dry regions in upper troposphere from north to south.
- Zones of low values indicate positions of ITCZ.





Difference of UTWV between 2002-2003 El Niño and 1999-2000 La Niña



UTWV Time Series (showing 30S – 30N)





Long-term UTWV Anomaly Dataset





UTWV Has Good Coverage for Tropical Convections



Conclusions

- Improved intersatellite calibration.
 - Temperature dependent inter-calibration
- Extension of time series to current.
 - HIRS/2 and HIRS/3 series connected
- The HIRS UTWV anomaly data are useful in monitoring Madden-Julian oscillation and various equatorial waves.



