

---

# Retrieval of Atmospheric Trace Gases Variability with Satellite Advanced IR Sounders

Alexander Uspensky, Anatoly Trotsenko\*, Alexander Kukharsky

SRC PLANETA, ROSHYDROMET, Moscow, Russia

\*RRC Kurchatov Institute, Moscow, Russia

## Outline


- IASI-based retrieval of atmospheric CH<sub>4</sub>, N<sub>2</sub>O, and CO columns
- Detecting of CO<sub>2</sub> variations from AIRS/Aqua data
- Concluding remarks

---

# IASI-based retrieval of atmospheric CH<sub>4</sub>, N<sub>2</sub>O, and CO columns

## Methods overview

# Methods overview: MPP Functional design



## Methods overview: MPP Content design

The **Modular Prototype Processor (MPP)** is the integrated software application capable of a self-contained execution of the procedures to retrieve the geophysical parameters from IASI level 1c data. MPP at the moment provides the following retrievals :

- land or sea surface skin temperature (SST)
  - temperature profile (0-40 km with 1 km resolution)
  - H2O total columnar amount
  - H2O mix. ratio profile (0-10 km with 1 km resolution)
  - O3 total columnar amount and partial column in between 0-20 km
  - O3 mix. ratio profile (20-40 km with 1 km resolution)
  - Minor gases total columns:
    - CH<sub>4</sub>
    - N<sub>2</sub>O
    - CO
- exploits
- measurements only
  - measurements only
  - SST and TP retrievals
  - SST and TP retrievals
  - SST and TP retrievals
  - SST and TP retrievals
  - SST and TP retrievals
  - Retrievals for:
    - SST, TP, H2O
    - SST, TP, H2O, CH4
    - SST, TP, H2O, O3


## MPP overview: Target characteristics

MPP retrievals	Target accuracy (r.m.s, per pixel)	Interfering factors to be a priori specified	A priori accuracy for interfering factors
Surface Skin Temperature	< 0.5-1.0 K	NO	-
Temperature profile	< 1-2K / 1 km	NO	-
H2O total column	< 10 %	SST TP	< 1 K < 2K /km
H2O mixing ratio profile (troposphere)	< 20-30% / 2km		
O3 total column	< 10 %	SST TP	< 1 K < 2K /km
O3 column 0-20 km	< 20 %		
O3 mixing ratio profile (20-40 km)	< 10 % / 1km		
CH4 total column	< 5 %	SST TP H2O (troposphere)	< 2 K < 2K /km < 30 % /2 km
N2O total column	< 10 %	SST TP H2O (troposphere) CH4 CA	< 2 K < 2K /km < 30 % /2 km < 10 %
CO total column	< 10 %	SST TP H2O (troposphere) O3 CA	< 2 K < 2K /km < 30 % /2 km < 10 %

## Methods overview: MPP approaches selection

Parameter	Retrieval Method
Surface Skin Temperature	Physical inversion, no online RTM use
Temperature profile	Regression-type, "ridge" scheme
H <sub>2</sub> O total column & mix. ratio profile	Regression-type, PCR technique, direct SST and TP use
O <sub>3</sub> total column & mix. ratio profile	1 <sup>st</sup> -Stage: Regression-type, PCR technique, direct SST and TP use
	2 <sup>nd</sup> -Stage: Physical inversion (15channels), Online RT modeling
CH <sub>4</sub> total column	Physical inversion, Online modeling (4 channels)
N <sub>2</sub> O total column	Physical inversion, Online RT modeling (3 channels)
CO total column	Physical inversion, Online RT modeling (5 channels)

# Methods overview: Channels



---

## Retrieval of Atmospheric Trace Gases Variability with IASI

Achieved retrieval accuracies




# MPP ACCURACIES: CH<sub>4</sub> column amount (Clear Sky)

Range of CH<sub>4</sub> column variation: from - 10% to +20 %

Sufficient accuracy (in terms of RMS) for supplementary T-q profiles retrieval : 2K and 30%, resp.

Sufficient accuracy (in terms of RMS) for supplementary N<sub>2</sub>O column retrieval: no requirement (!)

Retrieval error, %  
(in terms of rms)




# MPP ACCURACIES: N2O column amount (Clear Sky)

Range of N2O column variation: from - 10% to +20 %

Sufficient accuracy (in terms of RMS) for supplementary **T-q profiles** retrieval : **2K** and **30%**, resp.

Sufficient accuracy (in terms of RMS) for supplementary **CH4** column retrieval: **5-10 %**

Retrieval error, %



## CH<sub>4</sub> column amount retrieval from IASI L1c measurements

1. The CH<sub>4</sub> retrieval is carried out using **4** universal “static” IASI channels (i.e. independent on the lat/long zone and/or season): **1332.50, 1341.75, 1342.75 and 1346.75 cm<sup>-1</sup>**
2. The level of **retrieval accuracy** for the clear-sky conditions is at least **better than 5 %** (in terms of the r.m.s. error) for all seasons and lat/long zones, as well as rather wide range of the column variation (from **-10% to +30%**).
3. The above rather high accuracy is achieved **at moderate level of key interfering factors knowledge** correspondent to the r.m.s. accuracy (for the 1 km troposphere resolution) of **2K** and **30%** for the temperature and the water vapor, respectively

## N<sub>2</sub>O column amount retrieval from IASI L1c measurements

1. The prior estimate of the **CH<sub>4</sub>** with the accuracy at least not worse than **5%** provides to reliably retrieve **N<sub>2</sub>O** column in the clear-sky conditions with the r.m.s. accuracy ranging (for different lat/long zones and seasons) **from 4 to 9 %**
2. The level of the a priory knowledge about the temperature and humidity can be at the same level as for the methane case providing the retrieval of the **N<sub>2</sub>O** relative variations about + 10-20 %. For lesser levels of variations the retrieval accuracy is slightly degrades for several very cold and dry atmospheres **(to about 10%)**
3. Absolutely robust and reliable **N<sub>2</sub>O** retrieval (within **4-9 %** accuracy range) is available for all seasons and zones providing slightly better accuracy for the *a priory* T-q estimate, namely, **1.75 K** and **25%**, respectively.
4. The retrieval is reliably performed (in case of all lat/long zones and seasons) using **three IASI channels**, namely, those centered at **1277.25, 1298.50, 1299.50 cm<sup>-1</sup>**

# Effects of cloudiness in retrieving CH<sub>4</sub> and N<sub>2</sub>O from IASI level 1c measurements

## MOTIVATION:


1. Is it possible to retrieve trace gases columns from IASI measurements in case of partial or full cloud contaminated IFOV?
2. If it is possible to specify the limits for “applicable” cloudy situations and estimate the required accuracy level of the cloud parameters description

## EXPERIMENTS:

1. Retrieval of the CH<sub>4</sub> and N<sub>2</sub>O columns from the simulated IASI L1c measurements for characteristic sample of atmospheric situations (different lat/long zones, seasons, cloud coverage, cloud tops).
2. The retrievals have been performed in two “marginal” regimes:
  - a) the cloud parameters are specified accurately (referred to as “full cloud parameter correction”)
  - b) the cloud contamination of the IFOV is not taken into account in retrieving columns (referred to as “no cloud parameters correction”)

# CH<sub>4</sub>/N<sub>2</sub>O column amount retrieval from IASI: Effects of clouds

## General scheme of the retrieval experiment



# MPP ACCURACIES: CH<sub>4</sub> column amount (Effects of clouds)

ROOT MEAN SQUARE ERRORS (RMSE) of the CH<sub>4</sub> column amount retrieval (+20% shift),

Case: **Tropical latitudes**

Full cloud parameters correction

No cloud parameters correction

R.M.S. error , in %  
(Clear-sky case: 3.2 %)

Cloud Fraction

	0.1	0.2	0.3	0.5	0.7	1.0
1	3.2	3.2	3.2	3.2	3.2	3.2
2	3.2	3.2	3.2	3.2	3.1	3.2
3	3.2	3.2	3.2	3.3	3.3	3.5
5	3.2	3.3	3.4	3.3	4.4	5.4
7	3.3	3.5	3.8	5.3	6.4	9.1

Cloud Top , in km

R.M.S. error , in %  
(Clear-sky case: 3.2 %)

Cloud Fraction

	0.1	0.2	0.3	0.5	0.7	1.0
1	3.2	3.2	3.2	3.2	3.2	3.2
2	3.2	3.2	3.2	3.2	3.1	3.1
3	3.1	3.1	3.0	2.9	2.8	2.9
5	2.7	3.4	5.7	NOR	NOR	NOR
7	4.5	9.8	NOR	NOR	NOR	NOR

Cloud Top , in km

RMSE < 10 %

10% < RMSE < 20 %

RMSE > 20 % or No retrieval

# MPP ACCURACIES: N<sub>2</sub>O column amount (Effects of clouds)

ROOT MEAN SQUARE ERRORS (RMSE) of the N<sub>2</sub>O column amount retrieval (+20% shift),

Case: **Tropical latitudes**

Full cloud parameters correction

No cloud parameters correction

R.M.S. error , in %  
(Clear-sky case: 5.5 %)

Cloud Fraction

Cloud Top , in km	Cloud Fraction						
	0.1	0.2	0.3	0.5	0.7	1.0	
1	5.5	5.5	5.2	5.3	5.7	5.6	
2	5.2	5.2	5.2	5.7	6.0	6.9	
3	5.1	5.1	5.2	6.2	7.7	9.4	
5	5.4	5.7	6.1	8.2	10.3	15.6	
7	5.6	5.7	6.3	8.6	13.1	24.4	

R.M.S. error , in %  
(Clear-sky case: 5.5 %)

Cloud Fraction

Cloud Top , in km	Cloud Fraction						
	0.1	0.2	0.3	0.5	0.7	1.0	
1	5.2	5.4	5.9	6.6	7.1	8.2	
2	6.2	8.2	9.9	13.5	18.7	NOR	
3	7.1	10.0	13.3	20.4	NOR	NOR	
5	11.3	21.0	30.4	NOR	NOR	NOR	
7	18.8	NOR	NOR	NOR	NOR	NOR	

RMSE < 10 %

10% < RMSE < 20 %

RMSE > 20 % or No retrieval



## Trace gases retrieval: Conclusions

1. The retrieval of both the **CH<sub>4</sub> and N<sub>2</sub>O columns** is reliably possible for rather wide range of cloudy situations providing **high-accuracy knowledge** on key cloud (bulk) parameters such as the **cloud top** height and the **cover fraction**
2. Ignoring of the cloud contamination correction is feasible (i.e. provides reasonable column retrieval accuracy) **for the case of CH<sub>4</sub>** within the following cloud parameters limits:
  - H = 1 – 2 km ; fractions = 0.1 - 1.0
  - H = 3 – 5 km ; fractions = 0.1 - 0.3
  - H > 7 km ; fractions <= 0.1
3. Ignoring of clouds **for the case of N<sub>2</sub>O** retrieval is possible only for **low-level clouds (1-2 km) and low cover fractions (about 0.1 – 0.2)**
4. The **“feasibility limits” of cloud contamination ignoring** are valuably **wider** for sufficiently **wet and warm atmospheres** (tropics, middle latitude summer time) for both the CH<sub>4</sub> and N<sub>2</sub>O retrieval

# Towards estimating the column amounts of atmospheric CO<sub>2</sub> over boreal forests from AIRS/Aqua

- **CO<sub>2</sub> related issues**
- **Sensitivity studies**
- **AIRS channel selection for detecting CO<sub>2</sub> variations**
- **Retrieval of atmospheric CO<sub>2</sub> column amounts**
- **Validation exercise**

These studies have been financially supported by ESA/IAF (GMES Project “Development of novel techniques for CO<sub>2</sub> retrieval...”).


---

# CO2 related issues

**... “ consistent, very carefully calibrated studies of atmospheric CO2 are needed over long time periods to determine if, as suggested by data from recent year, CO2 growth rates and/or airborne fractions are increasing, or if there has been a shift in the land/ocean balance”**

(International Panel on Climate Change, 2006).


# Flask Sampling Networks. GlobalView 2000 (source: G.L.Stephens, R.Engelen)



# CO<sub>2</sub> in the atmosphere

CO<sub>2</sub> is no longer considered as uniformly mixed gas !

Instead, the goal is to determine small spatial regional variations of CO<sub>2</sub>




Annual cycle of CO<sub>2</sub> near-surface concentration BOREAS region (Canada). How far this variability goes in the atmosphere ?

Source:  
A.Trishchenko

# Satellite CO<sub>2</sub> observations (atmosphere)

- Solar region
  - SCIAMACHY/ENVISAT is the only sensor currently available**
  - Future Orbiting Carbon Observatory (OCO), GOSAT
  - New planned missions (Canadian GGAS-FTS, MEOS ...)
- Thermal region
  - AIRS/Aqua
  - Forthcoming IASI/METOP
  - some other sensors (IRFS-METEOR, CrIS/NPOESS, ...)



# Examples of CO<sub>2</sub> airborne observations



**Figure 3. Time series of CO<sub>2</sub> concentration at 1 km and 7 km altitudes over Surgut**

Note: Solid circles, blue lines, and green lines represent the observed data, the best-fit curves, and the long-term trends, respectively.

# Vertical structure and seasonal cycle of CO<sub>2</sub> over Siberian boreal zone



aircraft observations (Arshinov et al., 2005; Zuev et al., 2005)

Regional differences, amplitude of seasonal and diurnal cycle of CO<sub>2</sub> in the boundary layer can be very large (from 10-15 ppmv to 40-60 ppmv);  
Variability reduces with altitude, but still can be substantial (>5ppmv) up to the heights in the upper troposphere and over synoptic spatial scales.



# Critical Issues

**1. Retrieval of CO<sub>2</sub> column amount (CA) or profiles (CO<sub>2</sub>(p)) from atmospheric IR sounders data is problematic since the IR channels sensitive to CO<sub>2</sub> variations are also sensitive to temperature variations and the presence of clouds. Moreover the temperature and cloud variations should be treated as the main interfering factors. To advance the remote sensing methodology for monitoring CO<sub>2</sub> concentration from advanced IR sounders data means separating these effects.**

**2. To address this issue the studies have been focused on the selection of sub-sets of the dedicated AIRS super-channels. The objective of super-channel sub-sets selection is to reduce the effect of temperature profile uncertainties (key interfering factor) on the accuracy of trace gas CA retrieval. The signal in the dedicated super-channel (constructed as linear combination of pre-selected individual channels) should be sensitive to trace gas CA variations and should have small sensitivity to  $T(p)$  variations.**

# Procedures of AIRS data inversion and estimating CO<sub>2</sub> column amounts

**Two approaches are appears to be suitable for AIRS data inversion:**

- concurrent retrieval of “full” state vectors (incorporating some parameters relating to CO<sub>2</sub> abundance);**
  - self-dependent (or stand alone) retrieval of CO<sub>2</sub> abundance characteristics (using ancillary information, e. g. extracted from the same AIRS data).**
- The second approach has the advantage that it should be more flexible and less complicated.**

# Methodology

- **The approach to CO<sub>2</sub> concentration detection includes:**
- **collecting a dataset with representative samples of atmospheric state vectors and high resolution thermal infrared satellite measurements (AIRS) – these may be real or simulated via an adjusted Fast Radiative Transfer Models;**
- **analysis of the satellite measurement information content with respect to CO<sub>2</sub> perturbations and selection of dedicated AIRS channel subsets (to minimize interference from temperature and cloud effects);**
- **developing, testing, and validating the procedure for retrieval carbon dioxide CA (using synthetic or real satellite data).**

# Initial dataset


- **To generate a simulation dataset, one needs a representative sample of atmospheric state vectors that include the associated CO<sub>2</sub> CAs and/or vertical profiles. Various datasets can be utilized for compiling sample atmospheric state vectors, in particular, UKMO analysis data. In our studies we used two typical CO<sub>2</sub> vertical profiles extracted from (Schmidt et al., 1991) and complemented below 600 hPa by data from other sources. Some calculations and experiments have been performed using carbon dioxide profiles with constant mixing ratios and CAs (in the range of 350-380 ppmv).**
- **A sample of in-situ airborne CO<sub>2</sub> observations (over the area of boreal forests-Novosibirsk region as well as over Surgut region) was utilized in the validation exercise.**
- **The maximum CO<sub>2</sub> seasonal variability (about 16 ppmv) is expected at northern mid-latitudes between April and August. The variations of CO<sub>2</sub> CA in the range 5-15 ppmv were chosen in our studies.**

---

## **Fast Radiative Transfer Model for simulating AIRS data**


**Upon investigating SARTA (Stand alone AIRS Radiative Transfer Algorithm) from (Strow et al., 2003), we found the option to adjust CO<sub>2</sub>. At CIMSS the SARTA code has been implemented and modified by Yu. Plokhenko to accommodate changing CO<sub>2</sub> profiles. Later on the modified version of SARTA has been implemented at SRC “Planeta” (on a Windows platform) and used for modeling AIRS measurements and for conducting sensitivity studies.**

# Sensitivity Studies




The AIRS spectrum (between 500 and 3000 cm<sup>-1</sup>) of brightness temperature changes calculated with SARTA in response to decrease in total CO<sub>2</sub> from 370 ppmv to 350 ppmv (blue), to 355 ppmv (red), to 360 ppmv (black), and to 365 ppmv (green). The linearity of the response is evident.

**Variations of AIRS spectra due to CO<sub>2</sub> perturbations  
(optimal channel subset, 664-745 cm<sup>-1</sup>)**




**Variations of AIRS spectra due to CO<sub>2</sub> perturbations  
(optimal channel subset, 2249-2392 cm<sup>-1</sup>)**






**Variations of AIRS spectra due to CO2 perturbations  
(optimal channel subset, 664-745 cm-1)**



**Variations of AIRS spectra due to CO2 perturbations  
(optimal channel subset, 2249-2392 cm-1)**



# Approach to retrieval CO2 CA

- **It is necessary to specify channels with identical (or almost identical) and strongly differing response to T(p) and CO2 variations respectively (temperature- and CO2-dedicated).**

# Building of super-channels

- 1) Selection of T- and CO<sub>2</sub> -dedicated channels with “similar” temperature Jacobians or weighting functions HT; signals in one T-and several CO<sub>2</sub>-dedicated channels are designated as TB(I) and TB(II), TB(III),... respectively.
- 2) Specification of “synthetic” channel with signal TB(synth) as linear combination of two or more CO<sub>2</sub>-dedicated channels (signals TB(II), TB(III), TB(IV),...) and with temperature Jacobian, close to the Jacobian of T-dedicated channel. Signal in the synthetic channel is formed using least square fitting:  $TB(\text{synth}) = K_1 TB(\text{II}) + K_2 TB(\text{III})$ , where K<sub>1</sub>, K<sub>2</sub> are derived as solution of extremal problem:

$$\min J_w(K_1, K_2) = \sum_i [H_{T.I}(p_i) - H_{T.\text{synth}}(p_i)]^2, \quad H_{T.\text{synth}}(p) = K_1 H_{T.II}(p) + K_2 H_{T.III}(p).$$

- 3) Derivation of super-channel with signal  $TB(\text{sc}) = TB(I) - TB(\text{synth})$ .

# Building of super-channels

$$TB(sc) = TB.0(sc) + \delta T TB(sc) + \delta q TB(sc) + \delta Oz TB(sc) + TB(sc) + \varepsilon, \quad (1)$$

where:

$$\delta T TB(sc) = (HT.I - HT.synth) \Delta T,$$

$$\delta q TB(sc) = (Hq.I - Hq.synth) \Delta q,$$

$$\delta Oz TB(sc) = (HOz.I - HOz.synth) \cdot \Delta Oz,$$

$$TB(sc) = (HQ.I - HQ.synth) \Delta QCA,$$

TB.0(sc) is modeled signal calculated for vector  $\mathbf{x0}$ ; the temperature and other Jacobians are also calculated for  $\mathbf{x} = \mathbf{x0}$ .

Terms  $\delta T TB(sc)$ ,  $\delta q TB(sc)$ ,  $\delta Oz TB(sc)$  from (1) present input of variations T(p) and other interfering factors into signal variations  $\Delta TB(sc) = TB(sc) - TB.0(sc)$ .

**Variations  $|TB(sc)|$  should exceed notably variations  $|\delta T TB(sc)|$  as well as should exceed the instrumental noise.**

# METHOD OF AIRS DATA INVERSION

Formulae (\*) can be presented as follows:

$$\Delta TB(sc) = \kappa_1 + \kappa_2 \Delta QCA + \varepsilon,$$

$$\text{where } \kappa_1 = \delta T TB(sc) + \delta q TB(sc) + TB(sc), \kappa_2 = HQ(sc)$$

Linear relationship between  $\Delta QCA$  and measurements (variations  $\Delta TB(sc)$  or  $\Delta(sc)$  ) leads to the following regression estimator for  $\Delta QCA$  :

$$\Delta QCA(\text{regr}) = C_1 \Delta TB(sc) + C_2 ,$$

where  $C_1, C_2$  – const are regression coefficients.


There are two options:

- **Synthetic regression**-training sample is compiled using modeled AIRS data for various  $CO_2$  CA-s
- **Empirical regression**- training sample is compiled using collocated actual AIRS data and *in-situ* (airborne)  $CO_2$  observations


# SUPER-CHANNEL FORMATION FOR CO<sub>2</sub> CA RETRIEVAL

- **T-dedicated channel : #1917 at 2229.6 cm<sup>-1</sup>;**
- **CO<sub>2</sub>-dedicated channels: #2109 at 2388.2 cm<sup>-1</sup> and # 2113 at 2392.1 cm<sup>-1</sup>;**
- **TB(synth)=0.498 TB(2109) +0.426 TB(2113)**
- **TB(sc) = TB(1917) – TB(synth).**


Temperature Jacobians for T-, CO2-dedicated, and synthetic channels (Novosibirsk 3.3.2003)




Temperature Jacobians for T-, CO2-dedicated, and synthetic channels (Novosibirsk 17.9.2003)



Temperature Jacobians for T-, CO2-dedicated, and synthetic channels (Surgut 3.3.2003)




Temperature Jacobians for T-, CO2-dedicated, and synthetic channels (Surgut 17.9.2003)





# Synthetic Regression



**Error statistics for  $Q_{CA}$  retrieval  
(synthetic AIRS measurements, 2500 implementations)**


ppmv	360	365	370	375	380
<b>BIAS</b>	<b>0,06</b>	<b>0,04</b>	<b>-0,04</b>	<b>0,11</b>	<b>0,14</b>
<b>R.M.S.E.</b>	<b>2,11</b>	<b>2,07</b>	<b>1,98</b>	<b>2,00</b>	<b>2,03</b>

## Validation of the AIRS-based mid-tropospheric CO<sub>2</sub> CA retrievals over Siberian boreal forests





The series of retrieval experiments has been conducted for a sample of more than 10 granules of real AIRS measurements. It was found to be suitable to use the results of airborne measurements that have been performed within the framework of a joint Japanese-Russian Project, see e.g. (Arshinov et al., 2005). Regular measurements of the CO<sub>2</sub> concentration have been carried out since 1997 till now with the use of an Antonov-30 flying laboratory. The region of airborne surveys and the flight routes are located at the right bank of the southern part of the Ob Reservoir. The area of airborne measurements covers the region 54°08'-54°33' N, 81° 51'-82°40' E, moreover the boreal area consists 90% of coniferous trees. At the end of each month the ambient air is flask- sampled at heights of 0.5, 1, 1.5, 2, 3, 4, 5.5 and 7 km. Along with this the similar experiments have been conducted for the Surgut region (60-62° N, 70-75° E).

# Validation exercise for Novosibirsk region

Real AIRS data, namely granules with Cloud-cleared radiances for pre-selected area and time period between January and December 2003 (for one-two dates of each month) have been downloaded through [http://daac.gsfc.nasa.gov/data/dataset/AIRS/02\\_L2\\_Products/index.html](http://daac.gsfc.nasa.gov/data/dataset/AIRS/02_L2_Products/index.html). The atmospheric T- and q-profiles (components of state vectors) for the same area and time period have been extracted as already mentioned from the UKMO (Bracknell) analysis and complemented by results of ~~airborne CO2 measurements~~



**Flight route over region of interest**




Display (in HYDRA) of AIRS radiances in channels at 917.2, 2229.6, 2388.2, 2392.1  $\text{cm}^{-1}$  for the Novosibirsk region, 25 June 2003


# Novosibirsk region




# Results of validation exercise (actual AIRS data)



# Surgut region



# Results of validation exercise (actual AIRS data)





# Concluding remarks

The following developments are proposed to improve retrievals:

- **The creation of a data set of CO<sub>2</sub> mixing ratio profiles (from in situ airborne observations) representative of actual variability for different regions over boreal forests and complemented by co-located satellite measurements (AIRS, IASI) and atmospheric state vectors;**
- **The development and testing of the empirical and the synthetic regression estimators;**
- **The examination of the behavior of retrieval errors as a function of regression type, season, geography, and the number of pixels to be averaged in space and time (while applying the regression algorithm);**
- **The refinement and extension of the regression algorithm of CO<sub>2</sub> retrieval through more representative validation (involving more complete training/testing samples). Using the regression estimate as the first guess in the method of the retrieval of “full state vector”.**

# Selected References

- Aumann H.H. et al. *Geophys Res. Lett.*, 2005,32,L05806.
- Chédin A. et al. *JGR*, v. 108, N D2, 4064, 2003.
- Crevoisier C. et al. *Q.J.R. Meteorol. Soc.* v. 129. P.2719-2740, 2003.
- Rodgers C.D. *Proc. SPIE*, v. 283, P. 136-147, 1996.
- Strow L.L. et al. *IEEE Trans. Geosci. Rem. Sens.* v.41, N 2, P.303-313, 2003.
- Uspensky A.B., Kukharsky A.V., Trotsenko A.N. et al. “The 2005 Eumetsat Meteorol. Sat. Conf.” Dubrovnik, Croatia, P. 507-515, 2005.
- Uspensky A.B., Trotsenko A.N., Rublev A.N. *Earth Res. From Space*, N5, P.18-33, 2005 (in Russian).
- Uspensky A.B., Kukharsky A.V., Rublev A.N. *Earth Res. From Space*, N4, 2006, p. 42-51.
- Uspensky A.B., A.N. Trotsenko. *Proc. 1-st EPS/MetOp RAO Workshop. 15-17 May 2006, ESRIN, Frascati, Italy (ESA, SP-618, D.Danesy (Ed))*.
- Arshinov M. Yu., B.D.Belan, G.Inoue et al. In *Advances in the geological storage of carbone dioxide. International Approaches to reduce anthropogenic greenhouse gas emissions*. Ed. S. Lombardi, I .K. Altunina, S.T. Beanbien. Kluwer Acad. Publ., P. 11-16. 2005.
- Zuev V.V., B.D. Belan, N.E. Zueva et al., *Atmos. Oceanic Opt.*, v.18, N 7, P. 555-557,2005.

---

**THANKS**