

National Aeronautics and  
Space Administration

**Jet Propulsion Laboratory**  
California Institute of Technology  
Pasadena, California

# Variability in Eleven Years of AIRS Version 6 Observations

**Eric J. Fetzer, Joao Teixeira, Thomas Pagano and  
Bjorn Lambrigtsen**

**Jet Propulsion Laboratory / California Institute of Technology**


**International TOVS Study Conference  
Jeju Island, Korea**

**31 March 2014**



National Aeronautics and  
Space Administration  
  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# Atmospheric Infrared Sounder on Aqua in the A-Train






National Aeronautics and  
Space Administration  
  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# AIRS Key Level 2 Products


## Clouds and Water Vapor Feedback

CO




Greenhouse Gas Forcing


Atmospheric Temperature




Cloud Properties




Ozone




Atmospheric Water Vapor




Dust




Methane




CO2



SO2



Emissivity






National Aeronautics and  
Space Administration  
  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# AIRS Supporting Research

Over 631 AIRS Peer Reviewed Publications Through Oct 2013






National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## The Strengths of AIRS

*Most pertain to CrIS and IASI*

- **High infrared spectral resolution and coverage**  
*=> highest vertical resolution from the IR.*
- **Information about temperature and water vapor profiles, trace gases, etc. obtained simultaneously.**
- **Global coverage.**
- **11+ years of data (10 billion spectra, 1 billion retrievals).**



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## AIRS Challenges

- **In cloudy scenes most information is obtained in the microwave**  
⇒ *Lower vertical resolution than IR.*
- **Global coverage.**
- **11+ years of data (10 billion spectra, 1 billion retrievals).**




National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# Defining Tropical Conditions at 400 hPa: Potential Temperature > 310 K

6 Sep 2002



Level 3 Data




National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# Exploiting AIRS Strengths Relative Humidity at 400 hPa

6 Sep 2002

*Extremely demanding quality control  
( $<100\%$  yield for  $1 \times 1^\circ$  boxes in black).*





Level 3 Data






# 'Tropical' Conditions January 2003



**Occurrence Frequency,  $\theta > 310$  K at 400 hPa**

**Occurrence Frequency,  
Relative Humidity  $< 20\%$  at 400 hPa  
(NOT mean RH)**




National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# Defining 'Tropical' Conditions Dynamically


## July 2013

2013/07:  $\theta > 310$  K Occurrence Frequency (Percent),  $P = 400$  hPa



**Occurrence Frequency,  $\theta > 310$  K at 400 hPa**

2013/07: RH  $< 20\%$  Occurrence Frequency (Percent), Theta  $> 310$  K,  $P = 400$  hPa



**Occurrence Frequency,  
Relative Humidity  $< 20\%$  at 400 hPa  
(NOT mean RH)**




National Aeronautics and  
Space Administration  
  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California


# 400 hPa: Occurrence Frequency Weighted Area

$\theta > 310\text{ K}$  (thick)

$RH < 20\%$  (thin)

Their difference (dashed)





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Conclusions: Some Inside Information

- **AIRS has most information in clearer scenes**
  - *cloud-free conditions not required!*
- **Processes in the dry subtropics may be driving climate sensitivity. See:**
  - *Fasullo and Trenberth, 2013, Science.*
  - *Sherwood et al., 2014, Nature.*
- **With 11 years of observations, AIRS likely contains useful climate indices (like relative humidity quantities) in the dry tropics and subtropics.**
  - *Today's study is a preliminary attempt at creating one index.*