




# Impact Analysis of LEO Hyperspectral Sensor IFOV size on next generation NWP model forecast performance

Agnes Lim<sup>1</sup>, Allen Huang<sup>1</sup>, James Jung<sup>1</sup>, Zhenglong Li<sup>1</sup>, Jack Woollen<sup>2</sup>,  
FW Nagle<sup>1</sup>, Greg Quinn<sup>1</sup>, S. B. Healy<sup>3</sup>, Jason Otkin<sup>1</sup>, Mitch Goldberg<sup>4</sup> and Robert Atlas<sup>5</sup>

1. CIMSS/SSEC
2. IMSG/NOAA/NCEP/EMC
3. ECMWF
4. NOAA /JPSS Program Science Office
5. NOAA Atlantic Oceanographic and Meteorological Laboratory

4 Dec 2017


# Current CrIS FOVs verses next generation CrIS FOVs



# Current CrIS FOVs verses next generation CrIS FOVs

**Smaller FOV size → more chance of hole hunting for clear sky condition**

# Current CrIS FOVs verses next generation CrIS FOVs




Cloud Mask

# Current CrIS FOVs verses next generation CrIS FOVs


**Smaller FOV size → more chance of hole hunting for clear sky condition**

**How will assimilation of CrIS observations with smaller FOVs impact forecast performance?**

# Observing System Simulation Experiment




# Experiment Framework




# OSSE Calibration – Clear sky counts

Coverage for AIRS on  
2006-05-02 00z


Real



OSSE




OSSE  
tuned





# OSSE Calibration –QC count ratios



Ratio of used/rejected to total observation count  
considered for all used AIRS channels on 2006-05-02 00z

Real

OSSE

Tuned OSSE



# OSSE Calibration –

# Ratio of Standard Deviation


Before iterative tuning

After iterative tuning


RAOB uv




AIRS



ATMS




CrIS



N18 MHS

# Experiment Framework



## Impact assessment

- CTRL – CrIS 3x3, FOIV 5 @ 14km
- EXP – CrIS 6x6, all FOVs @ 7km

# Scorecard (2006-06-01 to 2006-06-30)

		P Lev	N. Hemisphere			S. Hemisphere			Tropics		
			Day 1	Day 3	Day 5	Day 1	Day 3	Day 5	Day 1	Day 3	Day 5
RMSE	Heights	100 hPa									
		500 hPa									
		850 hPa									
		1000 hPa									
	Wind Vector	200 hPa									
		500 hPa									
		850 hPa									
		1000 hPa									
	Temp	100 hPa									
		500 hPa									
850 hPa											
1000 hPa											
Bias	Heights	100 hPa									
		500 hPa									
		850 hPa									
		1000 hPa									
	Wind Vector	200 hPa									
		500 hPa									
		850 hPa									
		1000 hPa									
	Temp	100 hPa									
		500 hPa									
		850 hPa									
		1000 hPa									

<b>OSSE_EXP</b>		99.9% significant I
<b>better than</b>		99% significant level
<b>OSSE_CTRL</b>		95% significant level
Neutral		No statistical significant difference
<b>OSSE_CTRL</b>		95% significant level
<b>worse than</b>		99% significant level
<b>OSSE_EXP</b>		99.9% significant level

# Scorecard (2006-06-01 to 2006-06-30)


		P Lev	N. Hemisphere			S. Hemisphere		
			Day 1	Day 3	Day 5	Day 1	Day 3	Day 5
Anomaly Correlation	Heights	250 hPa						
		500 hPa						
		700 hPa						
		1000 hPa						
	Vector Wind	250 hPa						
		500 hPa						
		850 hPa						
	Temp	250 hPa						
		500 hPa						
		850 hPa						
MSLP	MSL							

<b>OSSE_EXP</b>	99.9% significant l
<b>better than</b>	99% significant level
<b>OSSE_CTRL</b>	95% significant level
Neutral	No statistical significant difference
<b>OSSE_CTRL</b>	95% significant level
<b>worse than</b>	99% significant level
<b>OSSE_EXP</b>	99.9% significant level


# OSSE deficiencies

- 1) Errors to be added to CrIS 6x6 observations
- 2) Calibration:
  - Statistics within 20%, issues with microwave mid tropospheric channels, infrared surface channels and tropopause.
  - Correlated errors
  - GPSRO error model
- 3) Observations simulation:
  - Conventional data
  - CrIS


# Comparison of CrIS simulation with different spatial resolution




CrIS 3x3 @ 14km



CrIS 6x6 @ 7km



# Number of clear observations per thinning box





# General deficiencies


## 1) Nature run

- Missing small scale resolvable features
- Cloud bias


## 2) GSI thinning

- One observation location per thinning box.

# Impact of different thinning resolution



# Impact of different thinning resolution



# Conclusion

- Forecast performance using higher spatial resolution CrIS observations
- A global OSSE system has been developed at CIMSS in coordination with NOAA OSSE team.
- Slight positive impact for bias, overall neutral results
- OSSE Deficiencies – simulation of observations, calibration and noise estimation for increased resolution CrIS.
- General Deficiencies – nature run and GSI thinning
- Better results expected with improvements in OSSE deficiencies .

**Under the support from JPSS Program Office, a global OSSE framework is developed, dedicated to support future JPSS impact studies.**

# Acknowledgements

We wish to thank:

- NASA GMAO for providing the Nature Run.
- Nigel Atkinson (UK Met Office) and Pascal Brunel (Meteo France) for providing information/test code for the modeling of unsteady non-zero yaw angles of METOP-A/B
- Radio Occultation Meteorology SAF for providing the Radio Occultation Processing Package as well as /technical support on the software usage.
- Sean Casey, Hongli Wang and Michiko Masutan for sharing their experience and providing various ancillary files needed by GFS.
- Szuchia Moeller, Nick Bearson and Derrick Herndon for post processing ATOVS data for orbit simulation comparison.
- Ron Errico for his advice on noise addition to simulated observations.
- Erik Olson for helping with computer resources for simulation of satellite radiances on SSEC Science Cluster.
- Sharon Nebuda for helping with the NCEP verifying package.
- NOAA/NESDIS. The experiments were run on the Supercomputer for Satellite Simulations and data assimilation Studies (S4) located at the University of Wisconsin– Madison