


Development of 3D Variational Assimilation System for ATOVS Data in China

Xue Jishan, Zhang Hua, Zhu Guofu, Zhuang Shiyu ¹⁾

Zhang Wenjian, Liu Zhiquan, Wu Xuebao, Zhang Fenyin.²⁾

¹⁾ Chinese Academy of Meteorological Sciences

²⁾ National Satellite Meteorological Center CMA




Contents


- *1. Introduction*
- *2. Variational data assimilation*
- *3. Assimilation of satellite radiances*
- *4. Impact of ATOVS data on typhoon prediction*
- *5. Towards operational implementation*

1. Introduction


Sparseness of observational data – the biggest challenge in improvement of weather forecasts



TEMP COVERAGE : /2003



The assimilation of satellite observations is in the first priority in the development of next generation NWP system




1. Introduction (cont.)

Two parallel projects for the application of TOVS data :

*Direct Assimilation within 3DVar frame work for
NWP in National Centers*

*Application of Retrieved Atmospheric Profiles to Local
NWP and Nowcasting*




Direct Assimilation of ATOVS Radiance

*A R&D project with joint efforts of scientists in
the National Satellite Meteorological Center
CMA and the Chinese Academy of
Meteorological Sciences*

Goal of the Project

*Alleviate the problem of data sparseness in
some crucial areas to which the prediction of
disastrous weather are sensitive*




Priorities :

Tropical Storms over Northwest Pacific


Regional Torrential Rains (usually caused by vortexes originating near the eastern wing of the Tibetan Plateau)

Studies focus at the application of ATOVS data to improve NWP of Typhoons (both track and intensity), especially those landing on Chinese coast



Direct Assimilation of ATOVS Radiance

as a sub-project of GRAPES: a 5-year project launched in 2001 aiming at the development of next generation numerical weather prediction in China



2. 3DVar in GRAPES

GRAPES: a NWP system newly developed for upgrading the operational medium range and mesoscale NWP's

Global / Regional Assimilation and Prediction System

Main Components of **GRAPES** :

Variational data assimilation

Unified nonhydrostatic model (grid mesh, SI/SL)

Model physics package

Parallel computing software



2. 3DVar in GRAPES (cont.)

Long / Lat Grid mesh

Control variables different from state variables


Preconditioning :

*With square root of back ground error covariance
matrix*

Recursive filter for limited area domain ;

Spectral filter for the globe

LBGFS for optimization



GRAPeS 3DVar

$$J = (X - X_b)^T B^{-1} (X - X_b) + (H(X) - Y_o)^T O^{-1} (H(X) - Y_o)$$

Analysis variables: Ψ, χ, T, q

Preconditioning with square root of
background error covariance matrix

Flexibility for different observational
operators

3. Direct Assimilation of Satellite Radiances in GRAPeS 3DVar

$$J = J_b + J_o$$


$$J_o = (Y - H(X))^T O^{-1} (Y - H(X))$$

$$H(X) : R * H * V(X)$$

R **Fast radiation transfer model (RTTOV is used)**

H **Horizontal interpolation**

V **Vertical Interpolation**




*Direct Assimilation of Satellite Radiances in **GRAPeS-3DVar***

Channel selection *General consideration :*

Channels sensitive to the surface characteristics, deep clouds and upper air (above 10 hpa) temperatures are avoided.


Noaa16/17 : AMSU-A CH 5-11

AMSU-B CH 18-20





Weighting functions (AMSU-A and B)

AMSU-A Channel Weighting Functions
(U.S. Standard Atmosphere)



Water vapor mixing ratio component weighting functions
(HIRS/3 channels 11 and 12, AMSU-B channels 1-5, U.S. standard atmosphere)





3. Assimilation of ATOVS data (cont.)


Acquisition and preprocessing of data (will be mentioned in 4)

Quality control before 3DVar :

Cloud identification

Bias correction

1DVar quality control




Two kinds of bias under consideration

- ⊕ Correction depending on scan angles: $s = \langle d_j(\theta) - d_j(\theta=0) \rangle$
- ⊕ Correction depending on air mass: $b = y - H(x_b) - s$
 - ⊠ Least square linear fitting
 - ⊠ Predictors p : air mass dependent

$$\mathbf{b} = \mathbf{A}\mathbf{p} + \mathbf{c}$$

$$\mathbf{A} = \mathbf{b}\mathbf{p}^T (\mathbf{p}\mathbf{p}^T)^{-1}$$



$$\mathbf{c} = \mathbf{b} - \mathbf{A}\mathbf{p}$$




Algorithms of bias correction


Following *Harris, Kelly(2001)*

- Scan angle correction- dependent on latitudes
- Predictors from the background:
 - Thickness between 1000-300hPa
 - Thickness between 200-50hPa
 - Surface temperatures
 - Integrated water vapor






Scan Bias




Scan Bias of NOAA16 AMSU-A CH 5-12 in the Zone 40N-50N

Samples for Statistics: Jul.1-10 2003 06/18 UTC (6 hours time window)




Comparison of NOAA16 AMSU-A before and after bias correction





Jul. 1-10 2003 06/18 UTC (6 hours time window)

Blue: before correction (background-obs.) ; Red: after correction;
Black: with standard deviation added and subtracted





NOAA16 AMSU-B






NOAA16 HIRS






AMSU-A Bias correction




Histogram of background-obs before bias correction July 15 2003 06UTC

Abscissa: TB (bin width 0.5deg) ; ordinate: number of obs within each bin




AMSU-A Bias correction




Histogram of background-obs after bias correction July 15 2003 06UTC

Abscissa: TB (bin width 0.5deg) ; ordinate: number of obs within each bin




AMSU-B before Bias correction




Histogram of background-obs before bias correction July 15 2003 06UTC

Abscissa: TB (bin width 0.5deg) ; ordinate: number of obs within each bin




AMSU-B after Bias correction




Histogram of background-obs after bias correction July 15 2003 06UTC

Abscissa: TB (bin width 0.5deg) ; ordinate: number of obs within each bin




HIRS before Bias correction




Histogram of background-obs before bias correction July 15 2003 06UTC

Abscissa: TB (bin width 0.5deg) ; ordinate: number of obs within each bin




HIRS after Bias correction



Histogram of background-obs after bias correction July 15 2003 06UTC

Abscissa: TB (bin width 0.5deg) ; ordinate: number of obs within each bin



4. Impact of ATOVS data on typhoon prediction

A case study :typhoon Rammason, June30-July6 2002

Data : radiosonde

ATOVS radiation (microwave)


background from T213 prediction

Prediction model : WRF



Control run: with only radiasondes

Exp1: with only ATOVS


Exp2: with ATOVS+ radiosondes



Coverage of ATOVS received by 3 ground stations





amsu-a (9600) , amsu-b (86670) , hirs (22400)



Impacts on forecast


*First Guess: 27 hours forecast by NMC's
T213 500hPa H*






Analysis of 500hPa H



Observations : Radiosondes 12UTC July 30 2002






Analysis of 500hPa H


Observations : ATOVS 17UTC July 30 2002







Analysis of 500hPa H


Observations : ATOVS 17UTC July 30 2002+
Radiosondes 12UTC July 30 2002






*First Guess: 27 hours forecast by NMC's
T213 500hPa q*






Analysis of 500hPa q



Observations : Radiosondes 12UTC July 30






Analysis of 500hPa q


Observations : ATOVS 17UTC June 30 2002
+radiosondes 12 UTC June30







Temperature deviation from zonal mean (along 23 N)

GRAPES_3DVAR BACKGROUND : SECTION of DT
DATE: 2002/07/02/18UTC




GRAPES_3DVAR ANALYSIS : SECTION of DT
DATE: 2002/07/02/18UTC








Zonal cross section of tangential winds (23N)

GRAPES_3DVAR BACKGROUND : SECTION of Vn
DATE: 2002/07/02/18UTC



GRAPES_3DVAR ANALYSIS : SECTION of Vn
DATE: 2002/07/02/18UTC







Analyses of moisture fields

GRAPES_3DVAR BACKGROUND : 850hPa Rh
DATE: 2002/07/02/18UTC


GRAPES_3DVAR ANALYSIS : 850hPa Rh
ANALYSIS DATE: 2002/07/02/18UTC




Background




With ATOVS



Analyses of moisture fields




GRAPES_3DVAR BACKGROUND : 600hPa Rh
DATE: 2002/07/02/18UTC





Background

GRAPES_3DVAR ANALYSIS : 600hPa Rh
ANALYSIS DATE: 2002/07/02/18UTC





With ATOVS




Analyses of moisture fields

GRAPES_3DVAR ANALYSIS : 400hPa Rh
ANALYSIS DATE: 2002/07/02/18UTC


GRAPES_3DVAR BACKGROUND : 400hPa Rh
DATE: 2002/07/02/18UTC



Background




With ATOVS





Vertical-zonal cross section of moisture

BACKGROUND : SECTION of RH
DATE: 2002/07/02/18UTC





GRAPES_3DVAR ANALYSIS : SECTION of RH
DATE: 2002/07/02/18UTC






Initial field: 850hpa H July 4 15UTC



Left: radiosondes

Right: radiosondes+ATOVS







Comparison between Predictions

Right : July 5 2002 12UTC 850 hPa H

Below: 21h prediction (radiosondes only)

Right below: 21h prediction (radiosondes+ATOVS)






Comparison between Predictions


Right: July 6 00UTC 850 hPa H (analysis)

Below: 33h prediction (radiosondes only) 探空资料, (35.5N,125.3E)


Right below: 33h prediction (radiosondes+ATOVS)




COLA/IGES 30 → 2002-12



COLA/IGES 30 → 2002-12



COLA/IGES 30 → 2002-12




Comparison between Predictions


Right: July 6 12UTC 850 hPa H (analysis)

Below: 45h prediction (radiosondes only)


Right below: 45h prediction (radiosondes +ATOVS)




COLA/IGES 30 2002-12-



COLA/IGES 30 2002-12-




COLA/IGES 40 2002-12-



Impact on the track prediction


TYPHOON TRACK




Starting from 15UTC July 4. 21,33,45 hours forecasts of the center's position are shown

5. Towards operational implementation

Flow Chart of Assimilation System on Pre-operational Trial





5. Towards operational implementation


Works in the near future :

Quality control

Usage of HIRS

AMSU over land

NOAA-16 AMSUB Brightness Temperature CH2
20020702 1702-1841-2023-2204UTC



160 180 200 220 240 260 280 300 320 340 360 380 400 K