



Progress and plans for the use of radiance data in the NCEP global and regional data assimilation systems



John Derber¹, Andrew Collard², Li Bi², Paul van Delst², Dave Groff², Jim Jung⁵, Xu Li², Emily H. C. Liu³,

Haixia Liu², Rahul Mahajan², Cathy Thomas², Russ Treadon¹, Xiaoyan Zhang⁴, Yanqiu Zhu²

¹NOAA/NCEP/EMC ²IMSG@NOAA/NCEP/EMC ³SRG@NOAA/NCEP/EMC ⁴ESSIC, University of Maryland

⁵University of Wisconsin

14th January 2015 Global Model Upgrade

- Add Metop-B IASI, F17 SSMIS,
- T1534 (~13km) forecast model.
- Analysis resolution remains on a linear grid corresponding to T574 (1152x576 grid boxes)
- Hybrid EnKF-3DVar data assimilation system is used with 75% of the solution prescribed by the ensemble and 25% from the static background error covariance.
- The resolution of the 80 member ensemble has increased from T254 to match the analysis resolution at T574.
- FASTEM-5
- Advanced radiance bias correction
- (including for SSMIS)

14th January 2015 Global Model Upgrade

- All-sky data assimilation for AMSU-A
- 4DEnsVar
- Temporal thinning.
- CRTM v2.2 including the FASTEM-6
- Assimilation of AVHRR and VIIRS AMVs
- Aircraft bias correction
- Monitor Saphir, GMI, AMSR-2 (JCSDA)

The future

- Extend All-sky data assimilation to other microwave and infrared instruments
- Cloud cleared radiances
- Spectrally correlated errors.
- FSO