

# Measurements of Stratospheric Volcanic Aerosol Optical Depth from NOAA/TOVS Observations



Clémence PIERANGELO<sup>1</sup>, Alain CHEDIN<sup>1</sup>, Patrick CHAZETTE<sup>2</sup>

<sup>1</sup>Laboratoire de Météorologie Dynamique, Institut Pierre-Simon Laplace – Ecole Polytechnique, 91128 Palaiseau cedex, FRANCE


<sup>2</sup>Laboratoire des Sciences du Climat et de l'Environnement, Institut Pierre-Simon Laplace – CEA, 91191 Gif sur Yvette Cedex, FRANCE




## Introduction :

In June 1991, the eruption of Mount Pinatubo (Philippines) injected about 20 Mt of sulfur dioxide into the stratosphere. The rapid conversion of sulfur dioxide into sulfuric acid (H<sub>2</sub>SO<sub>4</sub>) droplets caused a significant extinction, both in the visible and infrared. Here, we present a method to retrieve the Pinatubo aerosol optical depth at three infrared wavelengths (4.0 μm, 8.3 μm and 11.1 μm) from HIRS-2 measurements onboard NOAA-10 and NOAA-12 and collocated radiosonde measurements. No assumption on aerosol size distribution or refractive index is used, which is a major advantage of our method.


## 1. Signature of Pinatubo aerosol on CM-3 and CM-12 time series




## 3. Retrieval of aerosol optical depth



## 2. Simulation of the impact of volcanic aerosol on observations



## 4. Validation of the results



## Conclusions :

Covering the time period July 1987- September 1995, this analysis of differences between HIRS-2 measurements and collocated temperature and water vapor radiosonde profiles clearly reveals the signature of Mount Pinatubo aerosols. Simulations of their effect on brightness temperature using the climatological data base TIGR shows that, for a given channel, it depends on the atmospheric situation and linearly on the aerosol optical depth, and not on their microphysical properties. 3- or 12- month running mean aerosol optical depth at 4 μm, 8.3 μm and 11.1 μm are retrieved from channels 18, 10 and 8 measurements, for three latitude zone. Validation with lidar measurements and comparison with ISAMS measurements show good agreements.