

ECMWF is developing the capability to directly assimilate emitted clear-sky infrared limb radiances from the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS). It is the first time that the direct assimilation of limb radiances into a numerical weather prediction (NWP) system is being attempted. The developments have been prompted by the success of the direct assimilation of nadir radiances.

MIPAS is a very high spectral resolution sounder (0.025 cm⁻¹ wavenumber resolution) onboard ESA's Envisat satellite. It provides 59,605 spectral points or channels in 5 bands in the spectral range 685–2410 cm⁻¹. In the nominal scanning configuration, it provides one scan approximately every 5° latitude for 17 tangent altitudes between 6 and 68 km. The field of view (FOV) is 3 km in the vertical and 30 km in the horizontal.

Niels Bormann & Sean Healy

ECMWF, Shinfield Park,
Reading, Berkshire
RG2 9AX, UK

n.bormann@ecmwf.int



Niels Bormann

Channel selection and information content

A subset of MIPAS data has been selected for assimilation studies.

The iterative method of *Dudhia et al.* (2002) has been applied to select 325 channels over channel-specific tangent altitude ranges (Fig. 1).

This selection aims to maximise information content relative to the error estimate for the background data. The control variables are profiles of temperature, humidity, and ozone.

Figure 2 shows that MIPAS radiances have the potential to significantly reduce the analysis error of humidity and ozone throughout the stratosphere. Temperature errors are also significantly reduced in the stratosphere above about 30 km. MIPAS limb radiances are able to resolve background errors at much finer vertical resolution than nadir radiances (not shown).




Figure 1 Location of the selected MIPAS channels and tangent heights. Colour coding gives the ranking in the selection.




Figure 2 Theoretical reduction of analysis error that can be achieved with the selected MIPAS data. Black is the ECMWF background error; light blue the retrieval error expected if only MIPAS instrument noise and the background error are taken into account; orange is the estimate of errors introduced through the forward model (e.g., spectroscopic uncertainty, fixed climatology for certain gases, etc); blue the expected total retrieval error.

Assimilation framework

The above subset of MIPAS data is being used for assimilation studies with ECMWF's 4D-Var system.

- The radiative transfer model used is RTMIPAS, a regression-based fast radiative transfer model that follows RTTOV methodology (*Bormann et al.* 2004, 2005; see separate poster on RTMIPAS).
- Results presented here are based on using a 1-dimensional observation operator that assumes local horizontal homogeneity for the limb radiance calculations. Work on taking horizontal structure into account is in progress.
- Cloud-affected radiances are screened out using the method of *Remedios and Spang* (2002).
- Tangent pressure information is taken from ESA's level 2 data, since the satellite's engineering pointing information is not considered accurate enough.

Correction of radiance biases

Following the experience for nadir radiances, it is considered essential to correct so-called radiance biases in MIPAS data before the assimilation. If the MIPAS radiances are assimilated without bias correction, inconsistencies appear in the bias of analysis departures for different MIPAS radiances with weighting functions peaking at similar heights. Designing a bias correction for MIPAS radiances is made more difficult by the presence of biases in the model fields in the stratosphere, especially for humidity and ozone.

To resolve the ambiguity between model and radiance biases, an iterative method is being investigated, which involves repeated assimilation and tuning of subsets of the selected MIPAS channels over a 14-day study period. Such an iterative method provides some scope to separate between model and radiance biases, given that for each tangent altitude many MIPAS channels are available with weighting functions peaking around this tangent altitude.

Currently, bias correction is done with the so-called "gamma/delta" method (*Watts and McNally* 2004), which scales optical depths in the radiative transfer model with a channel-specific gamma, and models the remaining bias with a constant delta. This approach has been found to give a good model for the biases observed over a range of tangent altitudes for a large number of channels (Fig. 3).




Figure 3 FG bias (from 30 x 20° longitude/latitude boxes) vs. the signature of a 5% perturbation in gamma for a MIPAS temperature channel (both in nW/(cm² sr cm⁻¹)). Colour coding shows tangent height bands with grey for 60 km and green for 27 km.

Assimilation trials

Preliminary assimilation trials over the 14-day period 18–31 August 2003 show the following key results:

- MIPAS radiances can be assimilated while maintaining a similar level of fit to other observations sensitive to stratospheric temperature.
- The assimilation of MIPAS radiances has a significant impact on the mean temperature, humidity, and ozone analyses in the stratosphere, qualitatively in agreement with a reduction of known biases in the model fields (Fig. 4).
- Analyses with MIPAS radiances agree better with MIPAS level 2 retrievals in the regions expected from the information content study (Fig. 5). This provides a first cross-validation of the radiance assimilation.




Figure 4 a) Difference in mean temperature analyses [K] at 1 hPa between the experiment with and the one without MIPAS radiance assimilation. b) Relative difference in the mean humidity analyses [%] at 10 hPa between the experiment with and the one without MIPAS radiance assimilation.

Conclusions

Limb radiances from MIPAS have for the first time been directly assimilated into an NWP model. The results highlight the considerable potential of limb radiances to improve the analyses of temperature, humidity, and ozone in the stratosphere. Provided radiance biases are adequately addressed, assimilation of MIPAS radiances appears to correct considerable biases in the mean stratospheric analyses, and these biases qualitatively agree with known deficiencies in ECMWF model fields. Further work is needed to validate the resulting analyses against other observations in the stratosphere.

The above approach could be adopted to the assimilation of data from EOS-Aura's MLS.

References

- Bormann, N., M. Matricardi, and S.B. Healy**, 2005: A fast radiative transfer model for the assimilation of infrared limb radiances from MIPAS. *Quart. J. Roy. Meteor. Soc.*, **131**, in press.
- Bormann, N., M. Matricardi, and S.B. Healy**, 2004: RTMIPAS: A fast radiative transfer model for the assimilation of infrared limb radiances from MIPAS. *Tech. Memo. 436*, ECMWF, Reading, UK, 49 pp [available under www.ecmwf.int/publications/library/do/references/list/14].
- Dudhia, A., V.L. Jay, and C.D. Rodgers**, 2002: Microwindow selection for high-spectral-resolution sounders. *Applied Optics*, **41**, 3665–3673.
- Remedios, J., and R. Spang**, 2002: MIPAS observations of clouds and their effects on level 2 trace gas products. In *Envisat Validation Workshop*, Frascati, Italy, ESA, [available online under envisat.esa.int/pub/ESA_DOC/envisat_val_1202/proceedings/ACV/MIPAS/05_remedios.pdf].
- Watts, P.D., and A.P. McNally**, 2004: Identification and correction of radiative transfer model errors for atmospheric sounders: AIRS and AMSU-A. In *Proceedings of the ECMWF workshop on assimilation of high spectral resolution sounders in NWP*, Reading, UK, ECMWF, 23–38.




Figure 5 Departure statistics for ESA's MIPAS temperature retrievals (bias-solid; standard deviation-dotted) for the region 65–20°N, without (a) and with (b) MIPAS radiance assimilation.

Acknowledgements

Niels Bormann was funded through a shared-cost project co-funded by the Research Directorate General of the European Commission. Sean Healy is supported through the EUMETSAT/ECMWF Fellowship agreement. The ECMWF system has been developed by a large number of staff members and consultants at ECMWF and Météo France. All MIPAS data used © ESA.